ることができる。

以上の3種類の前葉体は概形，頂の彎入の状況，翼細胞の大きさ，造卵器の頸部の形状および頸細胞の数，造精器の大きさおよび生ずる場所等によつて相互に区別し得るが，総合的に見ると既報チャセンシダおよびトラノオシダとよく類似し，これらを含めた5種類の前葉体と同一型であるとみることができる。この型は *Asplenium* L. の基準種である A. *tricyomanes* L. を含む故，この型に属する種類をもつて前葉体からみた真正チャセンシダ属 *Euasplenium* を理解する。

○ナガサワハコベ追記（水島正美）　Masami MIZUSHIMA: Further note on the identity of *Stellaria laxa*

本誌32巻250～252頁(1957)において，台湾のナガサワハコベとフィリッピンの *S. laxa* Merr. とが東南アジア方面の *S. saxatilis* Buch.-Ham. に同じであることを論じた。其の際には *S. laxa* の基準標本を見ておらず，原記載に“種子が1.5～1.8 mm”とある点に多少の疑問を残しつゝ三者を合一したのであつた。其の後ニューヨーク植物園所蔵の副基準標本(Elmer 6612)を検する折に恵まれ，種子の長径が1.5mm或は微かに大きい程度であるのを確認した。故に三者が全く同一種に属することに疑う余地はない。

マニラにあつた正基準標本が戦災で失われた今日，ニューヨークにある同番号の副品を新基準標本と定めねばならない。此の標本に最も似た形のものは田川博士の no. 859 であり，台湾花蓮港庁の関ケ原～合歓での採集品である。熟した果実は見られないが十分参考になると思う。(東大理学部腊葉庫にある)。

In my discussion published in this journal (**32**: 250–252, 1957) on the identity of *Stellaria saxatilis* Buch.-Ham., *S. laxa* Merr. and *S. stellato-pilosa* Hayata, there remained a question to be settled. It is the problem of discrepancy in the size of seed at most 1.5 mm long against Merrill's description 1.5～1.8 mm long. Since I have not examined, at that time, the type of *S. laxa*, the reduction of two entities to *S. saxatilis* necessarily bear a slight hesitation. But now, through the courtesy of Dr. David D. Keck of the New York Botanical Garden, I was able to have a chance to examine the type collection of Merrill's species.

The specimen, *Elmer's 6612*, is a well grown plant and is labelled "*Stellaria laxa* Mrl., Co-type. Kias, Province of Benguet, Luzon, A. D. E. Elmer, June, 1904. From the Herbarium, Bureau of Government Laboratories, Manila, P. I.". The last abbreviation indicates apparently the Philippine Islands. The specimen has its sepals 3.5～4 mm long and free to the basal portion where is an obsolete receptacular thickening in fruit. There are a few capsules nearly ripe, and each

→ 246 p.

91) オオミ（大実）アセタケ（新種）。子実体は小形であるが胞子は大きい。夏秋季，大津市石山平津町，広葉樹下禾本草の間に発生する。

92) **ヤマノコガサ**。春—初夏，林内針葉樹の落葉，落枝，材の破片等に発生。大津市三井寺境内，奈良市春日神社附近で採つた。

93) **ケモミヴラモドキ**（新品種）。全体帯褐灰色で，微細な繊維状毛に密におおわれている。欧洲の *Rhodophyllus babingtonii* (Blox.) Quél. に比し傘がやや大きく，胞子の幅が広いのでその一品種とみなすことにした。京都大学植物園，および大津市近江神宮境内で採つた。

Errata

Journ. Jap. Bot. **33** (11): 344~350

Page	Line	for	read
344	25	29)	79)
	26	-pipe-	-pip-
346	5	conifers Bot.	conifers, Bot.
	31	pilea	pileo.
347	5	superior persistent	superior, persistent

Journ. Jap. Bot. **34** (7)

表紙裏　〔紙カットの説明〕→〔表紙カット……〕

221　表中 *D. crenata* (=*D. Sieboldi*) のカッコ内を 2 行下の *D. scabra* の行に移す。

---

238 p. →

contains a few seeds black-brown in colour. The seeds are 1.5 mm long or faintly more, but are less than 1.8 mm long. Thus the size of seeds does not go beyond that of *S. saxatilis* and *S. stellato-pilosa*. However, Merrill did describe the seeds as 1.5~1.8 mm long based on the type once kept in the Manila Herbarium in the Philippines and lost during the last war. Therefore the specimen Elmer's 6612, now preserved in the New York Herbarium, should be considered as neotype regardless the difference in the size of seed.

Finally, it is safely concluded that the Formosan *Stellaria stellato-pilosa* Hay. and the Philippine *S. laxa* Merr. are conspecific with the polymorphous southeastern Asiatic *S. saxatilis* Buch.-Ham. In my present knowledge, there is no geographical variety within the species. (Makino Herbarium, Tokyo Metropolitan University, Fukazawa-1., Setagaya, Tokyo. 東京都世田谷区深沢町，都立大学理学部牧野標本館)